

PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	<input checked="" type="checkbox"/> for					Go	Clear
Limits		Preview/Index		History		Clipboard	

Entrez PubMed

☐ 1: Dev Genet 1999;24(3-4):299-308[Related Articles, Books, LinkOut](#)

Peptides from the PKD repeats of polycystin, the PKD1 gene product, modulate pattern formation in the developing kidney.

van Adelsberg J.

Department of Medicine, Columbia University, New York, New York
10032, USA. jsv1@columbia.edu

Related Resources

Mutations in the PKD1 gene cause the majority of cases of autosomal dominant polycystic kidney disease. The PKD1 gene codes for a protein of unknown function, polycystin-1, that is predicted to be a receptor. Its large extracellular domain contains 16 copies of novel motif, the PKD repeat, that is likely to be a ligand binding domain based on its similarity to immunoglobulin domains. These observations suggested that soluble fragments of the extracellular domain of polycystin-1 could be used as competitive inhibitors of polycystin function in a suitable model system. Polycystin-1 is highly expressed in the ureteric bud and other branching epithelia during development and interacts with beta-catenin, a molecule known to play a role in branching morphogenesis. These data suggested that polycystin-1 might play a role in branching morphogenesis. I show here that peptides derived from the PKD repeats of polycystin-1 caused an asymmetric pattern of ureteric bud branching in cultured kidney rudiments. Treatment of kidney rudiments with experimental but not control peptides reduced both the number of ureteric bud branches and the number of nephrons. Experimental peptides produced significant morphogenetic effects at concentrations ≤ 0.1 mM. These data suggest that polycystin-1 plays a role in branching morphogenesis by the ureteric bud.

PMID: 10322638 [PubMed - indexed for MEDLINE]

Display	Abstract	<input checked="" type="checkbox"/>	Save	Text	Order	Add to Clipboard
---------	----------	-------------------------------------	------	------	-------	------------------



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	●MIM
Search PubMed	<input checked="" type="checkbox"/> for					Go	Clear
Limits		Preview/Index		History		Clipboard	
Display	Abstract	<input checked="" type="checkbox"/>	Save	Text	Order	Add to Clipboard	

Entrez PubMed

☐ 1: Lab Invest 1999 Oct;79(10):1311-23[Related Articles, Books, LinkOut](#)

PubMed Services

The PKD1 gene product, "polycystin-1," is a tyrosine-phosphorylated protein that colocalizes with alpha2beta1-integrin in focal clusters in adherent renal epithelia.


Wilson PD, Geng L, Li X, Burrow CR.

Department of Medicine, Mount Sinai School of Medicine, New York, New York 10029, USA. pat.wilson@SMTPlink.mssm.edu

Related Resources

Mutations in the PKD1 gene are responsible for autosomal dominant polycystic kidney disease (ADPKD). Although PKD1 has been cloned and shown to be expressed at high levels in the fetal ureteric bud and ADPKD cystic epithelia in the human kidney, the function of its encoded protein, "polycystin-1" is unknown. In this study we used primary and immortalized human renal epithelial cell lines derived from normal fetal, adult, and ADPKD kidneys, that endogenously express PKD1, to study the biologic function of the polycystin-1 protein. ADPKD renal epithelial cells expressed high levels of polycystin-1 protein and showed increased adhesion to type I collagen by comparison with normal adult human renal epithelia that expressed little polycystin. Adherent ADPKD cells also expressed high levels of alpha2beta1-integrin and their attachment was inhibited by a functional monoclonal antibody to alpha2-integrin. Double labeling and confocal microscopy as well as coimmunoprecipitation analysis showed overlapping colocalization of polycystin-1 with alpha2beta1-integrin as well as with the focal adhesion proteins vinculin and paxillin in multiprotein clusters localized to focal areas of cell membrane contact with type I collagen matrix after short periods of attachment. Immunoprecipitation and Western immunoblot studies also showed that polycystin-1 was posttranslationally modified by tyrosine phosphorylation. These studies suggest that the PKD1-encoded protein is part of a large multiprotein complex in epithelial cells that functions in the regulation of extracellular matrix interactions with the plasma membrane and cell cytoskeleton.

PMID: 10532593 [PubMed - indexed for MEDLINE]



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	<input checked="" type="checkbox"/> for					Go	Clear
Limits		Preview/Index		History		Clipboard	
Display	Summary	<input type="checkbox"/>	Save	Text	Order	Add to Clipboard	
Show: 20		<input type="checkbox"/> Items 1-20 of 168		Page 1 of 9		Select page: 1 2 3 4 5 6 7 8 9	

Entrez PubMed

PubMed Services

Related Resources


- ☐ **1:** [Li HP, Geng L, Burrow CR, Wilson PD.](#) Related Articles
Identification of phosphorylation sites in the PKD1-encoded protein C-terminal domain.
Biochem Biophys Res Commun. 1999 Jun 7;259(2):356-63.
PMID: 10362514 [PubMed - indexed for MEDLINE]
- ☐ **2:** [Parnell SC, Magenheimer BS, Maser RL, Calvet JP.](#) Related Articles
Identification of the major site of in vitro PKA phosphorylation in the polycystin-1 C-terminal cytosolic domain.
Biochem Biophys Res Commun. 1999 Jun 16;259(3):539-43.
PMID: 10364454 [PubMed - indexed for MEDLINE]
- ☐ **3:** [Sun G, Ke S, Budde RJ.](#) Related Articles
Csk phosphorylation and inactivation in vitro by the cAMP-dependent protein kinase.
Arch Biochem Biophys. 1997 Jul 15;343(2):194-200.
PMID: 9224730 [PubMed - indexed for MEDLINE]
- ☐ **4:** [Zamanillo D, Casanova E, Alonso-Llamazares A, Ovalle S, Chinchetru MA, Calvo P.](#) Related Articles, Protein, Nucleotide
Identification of a cyclic adenosine 3',5'-monophosphate-dependent protein kinase phosphorylation site in the carboxy terminal tail of human D1 dopamine receptor.
Neurosci Lett. 1995 Mar 31;188(3):183-6.
PMID: 7609904 [PubMed - indexed for MEDLINE]
- ☐ **5:** [Lee TR, Niu J, Lawrence DS.](#) Related Articles
The extraordinary active site substrate specificity of pp60c-src. A multiple specificity protein kinase.
J Biol Chem. 1995 Mar 10;270(10):5375-80.
PMID: 7534295 [PubMed - indexed for MEDLINE]
- ☐ **6:** [Saya H, Lee PS, Nishi T, Izawa I, Nakajima M, Gallick GE, Levin VA.](#) Related Articles
Bacterial expression of an active tyrosine kinase from a protein A/truncated c-src fusion protein.
FEBS Lett. 1993 Jul 26;327(2):224-30.
PMID: 7687570 [PubMed - indexed for MEDLINE]

- ☐ **7:** [Caohuy H, Pollard HB.](#) Related Articles
Activation of annexin 7 by protein kinase C in vitro and in vivo.
J Biol Chem. 2001 Apr 20;276(16):12813-21.
PMID: 11278415 [PubMed - indexed for MEDLINE]
- ☐ **8:** [Harada H, Becknell B, Wilm M, Mann M, Huang LJ, Taylor SS, Scott JD, Korsmeyer SJ.](#) Related Articles
Phosphorylation and inactivation of BAD by mitochondria-anchored protein kinase A.
Mol Cell. 1999 Apr;3(4):413-22.
PMID: 10230394 [PubMed - indexed for MEDLINE]
- ☐ **9:** [Wilson PD, Geng L, Li X, Burrow CR.](#) Related Articles
The PKD1 gene product, "polycystin-1," is a tyrosine-phosphorylated protein that colocalizes with alpha2beta1-integrin in focal clusters in adherent renal epithelia.
Lab Invest. 1999 Oct;79(10):1311-23.
PMID: 10532593 [PubMed - indexed for MEDLINE]
- ☐ **10:** [Izawa I, Tamaki N, Saya H.](#) Related Articles
Phosphorylation of neurofibromatosis type 1 gene product (neurofibromin) by cAMP-dependent protein kinase.
FEBS Lett. 1996 Mar 11;382(1-2):53-9.
PMID: 8612763 [PubMed - indexed for MEDLINE]
- ☐ **11:** [Sidovar MF, Kozlowski P, Lee JW, Collins MA, He Y, Graves LM.](#) Related Articles
Phosphorylation of serine 43 is not required for inhibition of c-Raf kinase by the cAMP-dependent protein kinase.
J Biol Chem. 2000 Sep 15;275(37):28688-94.
PMID: 10862777 [PubMed - indexed for MEDLINE]
- ☐ **12:** [Murphy BJ, Rogers J, Perdichizzi AP, Colvin AA, Catterall WA.](#) Related Articles
cAMP-dependent phosphorylation of two sites in the alpha subunit of the cardiac sodium channel.
J Biol Chem. 1996 Nov 15;271(46):28837-43.
PMID: 8910529 [PubMed - indexed for MEDLINE]
- ☐ **13:** [Briggs LJ, Stein D, Goltz J, Corrigan VC, Efthymiadis A, Hubner S, Jans DA.](#) Related Articles
The cAMP-dependent protein kinase site (Ser312) enhances dorsal nuclear import through facilitating nuclear localization sequence/importin interaction.
J Biol Chem. 1998 Aug 28;273(35):22745-52.
PMID: 9712906 [PubMed - indexed for MEDLINE]
- ☐ **14:** [Feschenko MS, Stevenson E, Sweadner KJ.](#) Related Articles
Interaction of protein kinase C and cAMP-dependent pathways in the phosphorylation of the Na,K-ATPase.
J Biol Chem. 2000 Nov 3;275(44):34693-700.
PMID: 10940309 [PubMed - indexed for MEDLINE]

- ☐ **15:** [Davare MA, Horne MC, Hell JW.](#) Related Articles
Protein phosphatase 2A is associated with class C L-type calcium channels (Cav1.2) and antagonizes channel phosphorylation by cAMP-dependent protein kinase.
J Biol Chem. 2000 Dec 15;275(50):39710-7.
PMID: 10984483 [PubMed - indexed for MEDLINE]
- ☐ **16:** [Reddy S, Aggarwal BB.](#) Related Articles
Curcumin is a non-competitive and selective inhibitor of phosphorylase kinase.
FEBS Lett. 1994 Mar 14;341(1):19-22.
PMID: 7511111 [PubMed - indexed for MEDLINE]
- ☐ **17:** [Park TS, Ostrander DB, Pappas A, Carman GM.](#) Related Articles
Identification of Ser424 as the protein kinase A phosphorylation site in CTP synthetase from *Saccharomyces cerevisiae*.
Biochemistry. 1999 Jul 6;38(27):8839-48.
PMID: 10393561 [PubMed - indexed for MEDLINE]
- ☐ **18:** [Sable CL, Filippa N, Hemmings B, Van Obberghen E.](#) Related Articles
cAMP stimulates protein kinase B in a Wortmannin-insensitive manner.
FEBS Lett. 1997 Jun 9;409(2):253-7.
PMID: 9202156 [PubMed - indexed for MEDLINE]
- ☐ **19:** [Arnould T, Kim E, Tsiokas L, Jochimsen F, Gruning W, Chang JD, Walz G.](#) Related Articles
The polycystic kidney disease 1 gene product mediates protein kinase C alpha-dependent and c-Jun N-terminal kinase-dependent activation of the transcription factor AP-1.
J Biol Chem. 1998 Mar 13;273(11):6013-8.
PMID: 9497315 [PubMed - indexed for MEDLINE]
- ☐ **20:** [Moyers JS, Zhu J, Kahn CR.](#) Related Articles
Effects of phosphorylation on function of the Rad GTPase.
Biochem J. 1998 Aug 1;333 (Pt 3):609-14.
PMID: 9677319 [PubMed - indexed for MEDLINE]

Display	Summary	Save	Text	Order	Add to Clipboard
Show: 20	<input checked="" type="checkbox"/> Items 1-20 of 168	Page 1 of 9		Select page: 1 2 3 4 5 6 7 8 9	

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM
Search PubMed	<input checked="" type="checkbox"/> for					Go	Clear
Limits		Preview/Index		History		Clipboard	

Entrez PubMed

Display	Summary	<input checked="" type="checkbox"/>	Save	Text	Order	Add to Clipboard				
Show: 20	<input checked="" type="checkbox"/>	Items 1-20 of 129	Page 1 of 7	Select page: 1	2	3	4	5	6	7

PubMed Services

- ☐ 1: [Wilson PD, Geng L, Li X, Burrow CR.](#) Related Articles

The PKD1 gene product, "polycystin-1," is a tyrosine-phosphorylated protein that colocalizes with alpha2beta1-integrin in focal clusters in adherent renal epithelia.

Lab Invest. 1999 Oct;79(10):1311-23.

PMID: 10532593 [PubMed - indexed for MEDLINE]

Related Resources

- ☐ 2: [Geng L, Burrow CR, Li HP, Wilson PD.](#) Related Articles

Modification of the composition of polycystin-1 multiprotein complexes by calcium and tyrosine phosphorylation.

Biochim Biophys Acta. 2000 Dec 15;1535(1):21-35.

PMID: 11113628 [PubMed - indexed for MEDLINE]

- ☐ 3: [Huan Y, van Adelsberg J.](#) Related Articles, OMIM

Polycystin-1, the PKD1 gene product, is in a complex containing E-cadherin and the catenins.

J Clin Invest. 1999 Nov;104(10):1459-68.

PMID: 10562308 [PubMed - indexed for MEDLINE]

- ☐ 4: [Geng L, Segal Y, Peissel B, Deng N, Pei Y, Carone F, Rennke HG, Glucksmann-Kuis AM, Schneider MC, Ericsson M, Reeders ST, Zhou J.](#) Related Articles

Identification and localization of polycystin, the PKD1 gene product.

J Clin Invest. 1996 Dec 15;98(12):2674-82.

PMID: 8981910 [PubMed - indexed for MEDLINE]

- ☐ 5: [van Adelsberg J.](#) Related Articles

Peptides from the PKD repeats of polycystin, the PKD1 gene product, modulate pattern formation in the developing kidney.

Dev Genet. 1999;24(3-4):299-308.

PMID: 10322638 [PubMed - indexed for MEDLINE]

- ☐ 6: [Ibraghimov-Beskrovnaya O, Bukanov NO, Donohue LC, Dackowski WR, Klinger KW, Landes GM.](#) Related Articles, OMIM

Strong homophilic interactions of the Ig-like domains of polycystin-1, the protein product of an autosomal dominant polycystic kidney disease gene, PKD1.

Hum Mol Genet. 2000 Jul 1;9(11):1641-9.

PMID: 10861291 [PubMed - indexed for MEDLINE]

- ☐ **7:** Van Adelsberg J, Chamberlain S, D'Agati V. Related Articles
Polycystin expression is temporally and spatially regulated during renal development.
Am J Physiol. 1997 May;272(5 Pt 2):F602-9.
PMID: 9176370 [PubMed - indexed for MEDLINE]
- ☐ **8:** Wilson PD, Burrow CR. Related Articles
Cystic diseases of the kidney: role of adhesion molecules in normal and abnormal tubulogenesis.
Exp Nephrol. 1999 Mar-Apr;7(2):114-24. Review.
PMID: 10213865 [PubMed - indexed for MEDLINE]
- ☐ **9:** Harris PC. Related Articles
Autosomal dominant polycystic kidney disease: clues to pathogenesis.
Hum Mol Genet. 1999;8(10):1861-6. Review.
PMID: 10469838 [PubMed - indexed for MEDLINE]
- ☐ **10:** Hanaoka K, Qian F, Boletta A, Bhunia AK, Piontek K, Tsiokas L, Sukhatme VP, Guggino WB, Germino GG. Related Articles, OMIM
Co-assembly of polycystin-1 and -2 produces unique cation-permeable currents.
Nature. 2000 Dec 21-28;408(6815):990-4.
PMID: 11140688 [PubMed - indexed for MEDLINE]
- ☐ **11:** van Adelsberg JS. Related Articles
The role of the polycystins in kidney development.
Pediatr Nephrol. 1999 Jun;13(5):454-9. Review.
PMID: 10412869 [PubMed - indexed for MEDLINE]
- ☐ **12:** Peters DJ, van de Wal A, Spruit L, Saris JJ, Breuning MH, Bruijn JA, de Heer E. Related Articles
Cellular localization and tissue distribution of polycystin-1.
J Pathol. 1999 Aug;188(4):439-46.
PMID: 10440756 [PubMed - indexed for MEDLINE]
- ☐ **13:** Palsson R, Sharma CP, Kim K, McLaughlin M, Brown D, Arnaout MA. Related Articles
Characterization and cell distribution of polycystin, the product of autosomal dominant polycystic kidney disease gene 1.
Mol Med. 1996 Nov;2(6):702-11.
PMID: 8972485 [PubMed - indexed for MEDLINE]
- ☐ **14:** Veldhuisen B, Spruit L, Dauwerse HG, Breuning MH, Peters DJ. Related Articles, Protein, Nucleotide, OMIM
Genes homologous to the autosomal dominant polycystic kidney disease genes (PKD1 and PKD2).
Eur J Hum Genet. 1999 Dec;7(8):860-72.
PMID: 10602361 [PubMed - indexed for MEDLINE]
- ☐ **15:** Geng L, Segal Y, Pavlova A, Barros EJ, Lohning C, Lu W, Nigam SK, Frischauf AM, Reeders ST, Zhou J. Related Articles
Distribution and developmentally regulated expression of murine polycystin.
Am J Physiol. 1997 Apr;272(4 Pt 2):F451-9.

PMID: 9140045 [PubMed - indexed for MEDLINE]

☐ 16: [Bateman A, Sandford R.](#)

[Related Articles](#)

The PLAT domain: a new piece in the PKD1 puzzle.
Curr Biol. 1999 Aug 26;9(16):R588-90. No abstract available.
PMID: 10469604 [PubMed - indexed for MEDLINE]

☐ 17: [Bycroft M, Bateman A, Clarke J, Hamill SJ, Sandford R, Thomas RL, Chothia C.](#)

[Related Articles](#), [Protein](#), [Structure](#)

The structure of a PKD domain from polycystin-1: implications for polycystic kidney disease.
EMBO J. 1999 Jan 15;18(2):297-305.
PMID: 9889186 [PubMed - indexed for MEDLINE]

☐ 18: [Ponting CP, Hofmann K, Bork P.](#)

[Related Articles](#)

A latrophilin/CL-1-like GPS domain in polycystin-1.
Curr Biol. 1999 Aug 26;9(16):R585-8. No abstract available.
PMID: 10469603 [PubMed - indexed for MEDLINE]

☐ 19: [Somlo S, Markowitz GS.](#)

[Related Articles](#)


The pathogenesis of autosomal dominant polycystic kidney disease: an update.
Curr Opin Nephrol Hypertens. 2000 Jul;9(4):385-94. Review.
PMID: 10926175 [PubMed - indexed for MEDLINE]

☐ 20: [Nauta J, Goedbloed MA, van den Ouweland AM, Nellist M, Hoogeveen AT.](#) [Related Articles](#)

Immunological detection of polycystin-1 in human kidney.
Histochem Cell Biol. 2000 Apr;113(4):303-11.
PMID: 10857482 [PubMed - indexed for MEDLINE]

Display	Summary	Save	Text	Order	Add to Clipboard
Show: 20	Items 1-20 of 129	Page 1 of 7	Select page: 1 2 3 4 5 6 7		

[Write to the Help Desk](#)
[NCBI](#) | [NLM](#) | [NIH](#)
[Department of Health & Human Services](#)
[Freedom of Information Act](#) | [Disclaimer](#)



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	
Search	PubMed	<input checked="" type="checkbox"/> for					Go	Clear
		Limits	Preview/Index	History	Clipboard			

Entrez PubMed

☐ 1: Exp Nephrol 1999 Mar-Apr;7(2):114-24 [Related Articles, Books, LinkOut](#)

PubMed Services

Cystic diseases of the kidney: role of adhesion molecules in normal and abnormal tubulogenesis.

Wilson PD, Burrow CR.

Mount Sinai School of Medicine, New York, N.Y., USA.
pat.wilson@SMTPlink.mssm.edu

Related Resources

This short review summarizes some information concerning what is known about matrix adhesion molecules, focal adhesion proteins, and cell-cell adhesion molecules in normal renal development and cystic diseases of the kidney. The focus is on human nephrogenesis and disease, but utilizes critical information gained from genetically manipulated mouse models. Interestingly, a significant role for the human PKD-1-encoded gene product, polycystin-1, has been found in cell-matrix interactions via integrins during development, and mutations lead to autosomal dominant polycystic kidney disease (ADPKD). Recent studies on human ADPKD have implicated polycystin-1 in the formation of multiprotein complexes containing focal adhesion proteins at the basal cell surface of the normal ureteric bud. Further evidence of a critical role of cell-matrix interactions via focal adhesion complex formation is provided by the development of renal cystic disease in tensin knockout mice.

Publication Types:


- Review
- Review, academic

PMID: 10213865 [PubMed - indexed for MEDLINE]

Display	Abstract	<input checked="" type="checkbox"/>	Save	Text	Order	Add to Clipboard
---------	----------	-------------------------------------	------	------	-------	------------------

[Write to the Help Desk](#)
NCBI | NLM | NIH

Department of Health & Human Services
Freedom of Information Act | Disclaimer



PubMed	Nucleotide	Protein	Genome	Structure	PopSet	Taxonomy	OMIM	
Search PubMed	<input checked="" type="checkbox"/> for						Go	Clear
Limits		Preview/Index		History		Clipboard		

Entrez PubMed

Display	Summary	<input checked="" type="checkbox"/>	Save	Text	Order	Add to Clipboard				
Show: 20	<input checked="" type="checkbox"/>	Items 1-20 of 137	Page 1 of 7	Select page: 1	2	3	4	5	6	7

PubMed Services

☐ 1: [van Adelsberg J.](#)

Related Articles

Peptides from the PKD repeats of polycystin, the PKD1 gene product, modulate pattern formation in the developing kidney.

Dev Genet. 1999;24(3-4):299-308.

PMID: 10322638 [PubMed - indexed for MEDLINE]

Related Resources

☐ 2: [Van Adelsberg J, Chamberlain S, D'Agati V.](#)

Related Articles

Polycystin expression is temporally and spatially regulated during renal development.

Am J Physiol. 1997 May;272(5 Pt 2):F602-9.

PMID: 9176370 [PubMed - indexed for MEDLINE]

☐ 3: [Bycroft M, Bateman A, Clarke J, Hamill SJ, Sandford R, Thomas RL, Chothia C.](#)

Related Articles, Protein, Structure

The structure of a PKD domain from polycystin-1: implications for polycystic kidney disease.

EMBO J. 1999 Jan 15;18(2):297-305.

PMID: 9889186 [PubMed - indexed for MEDLINE]

☐ 4: [Wilson PD, Geng L, Li X, Burrow CR.](#)

Related Articles

The PKD1 gene product, "polycystin-1," is a tyrosine-phosphorylated protein that colocalizes with alpha2beta1-integrin in focal clusters in adherent renal epithelia.

Lab Invest. 1999 Oct;79(10):1311-23.

PMID: 10532593 [PubMed - indexed for MEDLINE]

☐ 5: [Bateman A, Sandford R.](#)

Related Articles

The PLAT domain: a new piece in the PKD1 puzzle.

Curr Biol. 1999 Aug 26;9(16):R588-90. No abstract available.

PMID: 10469604 [PubMed - indexed for MEDLINE]

☐ 6: [Veldhuisen B, Spruit L, Dauwerse HG, Breuning MH, Peters DJ.](#)

Related Articles, Protein, Nucleotide, OMIM

Genes homologous to the autosomal dominant polycystic kidney disease genes (PKD1 and PKD2).

Eur J Hum Genet. 1999 Dec;7(8):860-72.

PMID: 10602361 [PubMed - indexed for MEDLINE]

- ☐ 7: Geng L, Segal Y, Pavlova A, Barros EJ, Lohning C, Lu W, Nigam SK, Frischauf AM, Reeders ST, Zhou J. Related Articles
Distribution and developmentally regulated expression of murine polycystin.
Am J Physiol. 1997 Apr;272(4 Pt 2):F451-9.
PMID: 9140045 [PubMed - indexed for MEDLINE]
- ☐ 8: Ponting CP, Hofmann K, Bork P. Related Articles
A latrophilin/CL-1-like GPS domain in polycystin-1.
Curr Biol. 1999 Aug 26;9(16):R585-8. No abstract available.
PMID: 10469603 [PubMed - indexed for MEDLINE]
- ☐ 9: Ward CJ, Turley H, Ong AC, Comley M, Biddolph S, Chetty R, Ratcliffe PJ, Gattner K, Harris PC. Related Articles, OMIM
Polycystin, the polycystic kidney disease 1 protein, is expressed by epithelial cells in fetal, adult, and polycystic kidney.
Proc Natl Acad Sci U S A. 1996 Feb 20;93(4):1524-8.
PMID: 8643665 [PubMed - indexed for MEDLINE]
- ☐ 10: Sandford R, Sgotto B, Aparicio S, Brenner S, Vaudin M, Wilson RK, Chisoe S, Pepin K, Bateman A, Chothia C, Hughes J, Harris P. Related Articles, Protein, Nucleotide
Comparative analysis of the polycystic kidney disease 1 (PKD1) gene reveals an integral membrane glycoprotein with multiple evolutionary conserved domains.
Hum Mol Genet. 1997 Sep;6(9):1483-9.
PMID: 9285785 [PubMed - indexed for MEDLINE]
- ☐ 11: Bogdanova N, McCluskey M, Sikmann K, Markoff A, Todorov V, Dimitrakov D, Schiavello T, Thomas M, Kalaydjieva L, Dworniczak B, Horst J. Related Articles
Screening the 3' region of the polycystic kidney disease 1 (PKD1) gene in 41 Bulgarian and Australian kindreds reveals a prevalence of protein truncating mutations.
Hum Mutat. 2000;16(2):166-74.
PMID: 10923038 [PubMed - indexed for MEDLINE]
- ☐ 12: Hughes J, Ward CJ, Peral B, Aspinwall R, Clark K, San Millan JL, Gamble V, Harris PC. Related Articles, Protein, Nucleotide, OMIM
The polycystic kidney disease 1 (PKD1) gene encodes a novel protein with multiple cell recognition domains.
Nat Genet. 1995 Jun;10(2):151-60.
PMID: 7663510 [PubMed - indexed for MEDLINE]
- ☐ 13: Palsson R, Sharma CP, Kim K, McLaughlin M, Brown D, Arnaout MA. Related Articles
Characterization and cell distribution of polycystin, the product of autosomal dominant polycystic kidney disease gene 1.
Mol Med. 1996 Nov;2(6):702-11.
PMID: 8972485 [PubMed - indexed for MEDLINE]
- ☐ 14: Peters DJ, Spruit L, Klingel R, Prins F, Baelde HJ, Giordano PC, Bernini LF, de Heer E, Breuning MH, Bruijn JA. Related Articles
Adult, fetal, and polycystic kidney expression of polycystin, the polycystic

kidney disease-1 gene product.

Lab Invest. 1996 Aug;75(2):221-30.

PMID: 8765322 [PubMed - indexed for MEDLINE]

- ☐ 15: [Kim E, Arnould T, Sellin L, Benzing T, Comella N, Kocher O, Tsiokas L, Sukhatme VP, Walz G.](#) **Free in PMC**, Related Articles, Protein, Nucleotide

Interaction between RGS7 and polycystin.

Proc Natl Acad Sci U S A. 1999 May 25;96(11):6371-6.

PMID: 10339594 [PubMed - indexed for MEDLINE]

- ☐ 16: [Thomas R, McConnell R, Whittacker J, Kirkpatrick P, Bradley J, Sandford R.](#) Related Articles, Protein, OMIM

Identification of mutations in the repeated part of the autosomal dominant polycystic kidney disease type 1 gene, PKD1, by long-range PCR.

Am J Hum Genet. 1999 Jul;65(1):39-49.

PMID: 10364515 [PubMed - indexed for MEDLINE]

- ☐ 17: [van Adelsberg JS.](#) Related Articles

The role of the polycystins in kidney development.

Pediatr Nephrol. 1999 Jun;13(5):454-9. Review.

PMID: 10412869 [PubMed - indexed for MEDLINE]

- ☐ 18: [Watnick T, Germino GG.](#) Related Articles

Molecular basis of autosomal dominant polycystic kidney disease.

Semin Nephrol. 1999 Jul;19(4):327-43. Review.

PMID: 10435671 [PubMed - indexed for MEDLINE]

- ☐ 19: [Parnell SC, Magenheimer BS, Maser RL, Rankin CA, Smine A, Okamoto T, Calvet JP.](#) Related Articles

The polycystic kidney disease-1 protein, polycystin-1, binds and activates heterotrimeric G-proteins in vitro.

Biochem Biophys Res Commun. 1998 Oct 20;251(2):625-31.

PMID: 9792824 [PubMed - indexed for MEDLINE]

- ☐ 20: [Nauta J, Goedbloed MA, van den Ouweland AM, Nellist M, Hoogeveen AT.](#) Related Articles

Immunological detection of polycystin-1 in human kidney.

Histochem Cell Biol. 2000 Apr;113(4):303-11.

PMID: 10857482 [PubMed - indexed for MEDLINE]

Display	Summary	<input checked="" type="checkbox"/>	Save	Text	Order	Add to Clipboard					
Show: 20	<input checked="" type="checkbox"/>	Items 1-20 of 137	Page 1 of 7	Select page:	1	2	3	4	5	6	7

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

09478737
***** STN Columbus *****

FILE 'HOME' ENTERED AT 08:08:46 ON 31 JUL 2001

=> file medline biosis embase caplus uspatfull

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'MEDLINE' ENTERED AT 08:08:58 ON 31 JUL 2001

FILE 'BIOSIS' ENTERED AT 08:08:58 ON 31 JUL 2001

COPYRIGHT (C) 2001 BIOSIS(R)

FILE 'EMBASE' ENTERED AT 08:08:58 ON 31 JUL 2001

COPYRIGHT (C) 2001 Elsevier Science B.V. All rights reserved.

FILE 'CAPLUS' ENTERED AT 08:08:58 ON 31 JUL 2001

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'USPATFULL' ENTERED AT 08:08:58 ON 31 JUL 2001

CA INDEXING COPYRIGHT (C) 2001 AMERICAN CHEMICAL SOCIETY (ACS)

=> s polycystin (p) screen (p) atpase (p) collagen (p) focal (p) adhesion

L1 0 POLYCYSTIN (P) SCREEN (P) ATPASE (P) COLLAGEN (P) FOCAL (P)
ADHESION

=> s polycystin (p) screen

L2 12 POLYCYSTIN (P) SCREEN

=> dup rem 12

PROCESSING COMPLETED FOR L2

L3 4 DUP REM L2 (8 DUPLICATES REMOVED)

=> d 13 total ibib kwic

L3 ANSWER 1 OF 4 BIOSIS COPYRIGHT 2001 BIOSIS

ACCESSION NUMBER: 2001:302612 BIOSIS

DOCUMENT NUMBER: PREV200100302612

TITLE: RGS14, a regulator of heterotrimeric G-protein signaling
localizes to centrosome and interacts with a centrosome
protein, ninien.

AUTHOR(S): Cho, Hyeseou (1); Kehrl, John H. (1)

CORPORATE SOURCE: (1) NIH, 10 Center Dr., Bldg. 10 Rm. 11B08, Bethesda, MD,
20892 USA

SOURCE: FASEB Journal, (March 7, 2001) Vol. 15, No. 4, pp. A577.
print.

American Meeting Info.: Annual Meeting of the Federation of
Societies for Experimental Biology on Experimental Biology

2001 Orlando, Florida, USA March 31-April 04, 2001
ISSN: 0892-6638.

DOCUMENT TYPE: Conference
LANGUAGE: English
SUMMARY LANGUAGE: English

AB. . . contain additional domains known to interact with a variety of signaling molecules such as 14-3-3, Rap1/2, RhoA, Gbeta5, GIPC, and **polycystin**. RGS14, a larger member of the RGS family impairs G1alpha- and G13alpha-mediated signaling pathways and is strongly expressed in lymphocytes. To search for additional RGS14 functions, we performed a yeast 2-hybrid **screen** using a human spleen library with an RGS14 bait. We identified the human centrosome protein, ninein as well as several. . .

L3 ANSWER 2 OF 4 MEDLINE DUPLICATE 1
ACCESSION NUMBER: 2001031117 MEDLINE
DOCUMENT NUMBER: 20490726 PubMed ID: 10913159
TITLE: In vivo interaction of the adapter protein CD2-associated protein with the type 2 polycystic kidney disease protein, polycystin-2.
AUTHOR: Lehtonen S; Ora A; Olkkonen V M; Geng L; Zerial M; Somlo S;
CORPORATE SOURCE: Lehtonen E
Department of Pathology, Haartman Institute, University of Helsinki, P. O. Box 21, FIN-00014 Helsinki, Finland.
CONTRACT NUMBER: P50DK57328 (NIDDK)
SOURCE: JOURNAL OF BIOLOGICAL CHEMISTRY, (2000 Oct 20) 275 (42) 32888-93.
Journal code: HIV. ISSN: 0021-9258.
PUB. COUNTRY: United States
Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 200011
ENTRY DATE: Entered STN: 20010322
Last Updated on STN: 20010322
Entered Medline: 20001120

AB . . . at lower levels in renal tubular epithelial cells in the adult kidney, particularly in distal nephron segments. Independent yeast two-hybrid **screens** using the COOH-terminal region of either CD2AP or **polycystin-2** as bait identified the COOH termini of **polycystin-2** and CD2AP, respectively, as strong interacting partners. This interaction was confirmed in cultured cells by co-immunoprecipitation of endogenous **polycystin-2** with endogenous CD2AP and vice versa. CD2AP shows a diffuse reticular cytoplasmic and perinuclear pattern of distribution, similar to **polycystin-2**, in cultured cells, and the two proteins co-localize by indirect double immunofluorescence microscopy. CD2AP is an adapter molecule that associates. . . membrane proteins to organize the cytoskeleton around a polarized site. Such a function fits well with that hypothesized for the **polycystin** proteins in renal tubular epithelial cells, and the present findings suggest that CD2AP has a role in **polycystin-2** function.

L3 ANSWER 3 OF 4 MEDLINE DUPLICATE 2
ACCESSION NUMBER: 1998104524 MEDLINE
DOCUMENT NUMBER: 98104524 PubMed ID: 9442442
TITLE: Autosomal dominant polycystic kidney disease: clinical and genetic aspects.
AUTHOR: Sessa A; Ghiggeri G M; Turco A E
CORPORATE SOURCE: Department of Nephrology, G. Gaslini Children's Hospital, Genova, Italy.
SOURCE: JOURNAL OF NEPHROLOGY, (1997 Nov-Dec) 10 (6) 295-310.
Ref: 184
Journal code: CWE; 9012268. ISSN: 1120-3625.

PUB. COUNTRY: Italy
Journal; Article; (JOURNAL ARTICLE)
General Review; (REVIEW)
(REVIEW, TUTORIAL)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 199802
ENTRY DATE: Entered STN: 19980306
Last Updated on STN: 19980306
Entered Medline: 19980226

AB . . . studies on cystogenesis suggest a key role of cell-to-cell or cell-to-matrix interactions. Surface proteins mediating cell-to-cell contact, such as E-cadherin (**polycystin**?), integrin interactions, growth factors, receptor expression, are involved in the process of differentiation of the cellular condition and of the . . . (unknown chromosome) in a few families. PCR-based mutation detection methods, automated DNA sequencing, and other "functional" methods are used to **screen** and analyse ADPKD patients. It is not yet known whether the mutations identified so far in PKD1 and PKD2 inactivate the genes or generate an aberrant product. The products of PKD1 and PKD2 genes have been called **polycystin** 1 and 2. **Polycystins** are members of a family of interactive proteins involved in complex adhesive cell-cell, cell-matrix, protein-protein, and protein-carbohydrate interactions in the. . .

L3 ANSWER 4 OF 4 MEDLINE DUPLICATE 3
ACCESSION NUMBER: 96108969 MEDLINE
DOCUMENT NUMBER: 96108969 PubMed ID: 8554072
TITLE: Screening the 3' region of the polycystic kidney disease 1 (PKD1) gene reveals six novel mutations.
AUTHOR: Peral B; San Millan J L; Ong A C; Gamble V; Ward C J; Strong C; Harris P C
CORPORATE SOURCE: MRC Molecular Haematology Unit, Institute of Molecular Medicine, John Radcliffe Hospital, Headington, Oxford, United Kingdom.
SOURCE: AMERICAN JOURNAL OF HUMAN GENETICS, (1996 Jan) 58 (1) 86-96.
Journal code: 3IM; 0370475. ISSN: 0002-9297.
PUB. COUNTRY: United States
Journal; Article; (JOURNAL ARTICLE)
LANGUAGE: English
FILE SEGMENT: Priority Journals
ENTRY MONTH: 199602
ENTRY DATE: Entered STN: 19960306
Last Updated on STN: 19960306
Entered Medline: 19960221

AB . . . kidney disease (ADPKD), PKD1 (polycystic kidney disease 1), has been fully characterized and shown to encode an integral membrane protein, **polycystin**, involved in cell-cell and/or cell-matrix interactions. Study of the PKD1 gene has been complicated because most of the gene lies. . . elsewhere on the same chromosome, and consequently only seven mutations have been described so far. Here we report a systematic **screen** covering approximately 80% of the approximately 2.75 kb of translated transcript that is encoded by single-copy DNA. We have identified. . . they indicate that the majority of mutations lie within the duplicated area, which is difficult to study. The regions of **polycystin** removed in each mutation so far described are assessed for their functional significance; an area disrupted by two new small. . .

=> s polycystin (p) atpase (p) collagen (p) focal (p) adhesion

L4 1 POLYCYSTIN (P) ATPASE (P) COLLAGEN (P) FOCAL (P) ADHESION

=> d l4 total kwic

L4 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2001 ACS

AB . . . to identify agents that regulate the activity of the polycystic kidney disease proteins encoded by the PKD-1 and PKD-2 genes (**polycystin**-1 and -2) and that may be useful in the treatment of polycystic kidney disease. The assays of the invention comprise. . . decrease in the PKD mediated mutant phenotype. Characteristics assocd. with such a mutant phenotype include increased adherence to type I **collagen**-coated surfaces; apical expression of NaK-**ATPase** on the cell membrane; increased expression of .beta.-2-NaK-**ATPase**; and decreased **focal adhesion** kinase (FAK) incorporation into **focal adhesion** complexes, and inability to form tubular structures in a gel matrix. To facilitate the screening methods of the invention, cells. . . engineered to express epitope tagged PKD gene products and/or epitope tagged PKD interacting proteins (PKD-IP). Such interacting proteins include e.g. **focal adhesion** complex proteins such as FAK, paxillin, vinculin, and talin.

=> d l4 total ibib kwic

L4 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 2001:507954 CAPLUS

TITLE: Polycystin-based screening methods for compounds useful in the treatment of polycystic kidney disease

INVENTOR(S): Wilson, Patricia D.; Burrow, Christopher R.

PATENT ASSIGNEE(S): Mount Sinai School of Medicine of New York University,

USA

SOURCE: PCT Int. Appl., 56 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001050130	A2	20010712	WO 2001-US100317	20010105
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
PRIORITY APPLN. INFO.:			US 2000-478737	A 20000106
			US 2000-689461	A 20001012

AB Cell-based screening assays are provided which are designed to identify agents that regulate the activity of the polycystic kidney disease proteins encoded by the PKD-1 and PKD-2 genes (**polycystin**-1 and -2) and that may be useful in the treatment of polycystic kidney disease. The assays of the invention comprise the contacting of genetically engineered cells expressing a mutant or truncated PKD gene product with a test agent and assaying for a decrease in the PKD mediated mutant phenotype. Characteristics assocd. with such a mutant phenotype include

increased adherence to type I **collagen**-coated surfaces; apical expression of NaK-**ATPase** on the cell membrane; increased expression of .beta.-2-NaK-**ATPase**; and decreased **focal adhesion** kinase (FAK) incorporation into **focal adhesion** complexes, and inability to form tubular structures in a gel matrix. To facilitate the screening methods of the invention, cells may be genetically engineered to express epitope tagged PKD gene products and/or epitope tagged PKD interacting proteins (PKD-IP). Such interacting proteins include e.g. **focal adhesion** complex proteins such as FAK, paxillin, vinculin, and talin.

=> s polycystin (p) assay (p) expression

L5 11 POLYCYSTIN (P) ASSAY (P) EXPRESSION

=> dup rem l5

PROCESSING COMPLETED FOR L5

L6 4 DUP REM L5 (7 DUPLICATES REMOVED)

=> d l6 total ibib kwic

L6 ANSWER 1 OF 4 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 2001:507954 CAPLUS

TITLE: Polycystin-based screening methods for compounds useful in the treatment of polycystic kidney disease

INVENTOR(S): Wilson, Patricia D.; Burrow, Christopher R.

PATENT ASSIGNEE(S): Mount Sinai School of Medicine of New York University,

USA

SOURCE: PCT Int. Appl., 56 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001050130	A2	20010712	WO 2001-US100317	20010105
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
PRIORITY APPLN. INFO.:			US 2000-478737	A 20000106
			US 2000-689461	A 20001012

AB Cell-based screening **assays** are provided which are designed to identify agents that regulate the activity of the polycystic kidney disease proteins encoded by the PKD-1 and PKD-2 genes (**polycystin** -1 and -2) and that may be useful in the treatment of polycystic kidney disease. The **assays** of the invention comprise the contacting of genetically engineered cells expressing a mutant or truncated PKD gene product with a test agent and assaying for a decrease in the PKD mediated mutant phenotype. Characteristics assocd. with such a mutant phenotype include increased adherence to type I collagen-coated surfaces; apical **expression** of NaK-ATPase on the cell membrane; increased **expression** of .beta.-2-NaK-ATPase; and decreased focal adhesion kinase (FAK) incorporation into focal adhesion complexes, and inability

to

form tubular structures in a gel matrix. To facilitate the screening methods of the invention, cells may be genetically engineered to express epitope tagged PKD gene products and/or epitope tagged PKD interacting proteins (PKD-IP). Such interacting proteins include e.g. focal adhesion complex proteins such as FAK, paxillin, vinculin, and talin.

L6 ANSWER 2 OF 4 MEDLINE DUPLICATE 1
 ACCESSION NUMBER: 2001017517 MEDLINE
 DOCUMENT NUMBER: 20380163 PubMed ID: 10926175
 TITLE: The pathogenesis of autosomal dominant polycystic kidney disease: an update.
 AUTHOR: Somlo S; Markowitz G S
 CORPORATE SOURCE: Department of Internal Medicine (Nephrology), Yale University School of Medicine, USA.
 CONTRACT NUMBER: DK54053 (NIDDK)
 DK57328 (NIDDK)
 SOURCE: CURRENT OPINION IN NEPHROLOGY AND HYPERTENSION, (2000 Jul) 9 (4) 385-94. Ref: 69
 Journal code: B4H. ISSN: 1062-4821.
 PUB. COUNTRY: ENGLAND: United Kingdom
 Journal; Article; (JOURNAL ARTICLE)
 General Review; (REVIEW)
 (REVIEW, TUTORIAL)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 200011
 ENTRY DATE: Entered STN: 20010322
 Last Updated on STN: 20010322
 Entered Medline: 20001107

AB . . . normal allele in individual polarized epithelial cells. Most recent advances are focused on the function of the respective protein products, **polycystin-1** and **polycystin-2**. Indirect evidence supports an interaction between **polycystin-1** and -2, albeit it is unlikely that they work in concert in all tissues and at all times. They associate in yeast two hybrid and cotransfection **assays** and there is a striking similarity in the renal and pancreatic cystic phenotypes of Pkd2-/- and Pkd1del34/del34 mice. Also, the . . . human disease phenotypes remain completely overlapping with the major difference being in relative severity. Mounting evidence supports the hypothesis that **polycystin-1** is a cell surface receptor. A close homologue in the sea urchin sperm mediates the acrosome reaction in response to. . . disease domains reveals a beta-sandwich fold commonly found in surface receptor molecules. Indirect evidence also supports the initial hypothesis that **polycystin-2** is a calcium channel subunit. Several closely related homologues retain the calcium channel signature motif but differ in their predicted. . . shown to be a calcium regulated cation channel. Several important distinctions in polcystin-1 and -2 function have also been discovered. **Polycystin** -2 has a role in cardiac development that polcystin-1 does not. High level

polycystin-2 expression in renal epithelial cells coincides with maturation and elongation of tubules and, unlike **polycystin-1**, persists into adulthood. In cells in tissue culture, **polycystin-2** is expressed exclusively in the endoplasmic reticulum whilst the cellular **expression** of **polycystin-1** remains unknown. Overall, the difficult task of understanding the autosomal dominant polycystic disease process is proceeding apace.

L6 ANSWER 3 OF 4 MEDLINE DUPLICATE 2
 ACCESSION NUMBER: 1999293063 MEDLINE
 DOCUMENT NUMBER: 99293063 PubMed ID: 10362514
 TITLE: Identification of phosphorylation sites in the PKD1-encoded
 protein C-terminal domain.
 AUTHOR: Li H P; Geng L; Burrow C R; Wilson P D
 CORPORATE SOURCE: Department of Medicine, Mount Sinai School of Medicine, New

CONTRACT NUMBER: York, New York 10029, USA.. Hsi-Ping_Li@mtplink.mssm.edu
 SOURCE: R01 DK448833 (NIDDK)
 BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, (1999
 Jun 7) 259 (2) 356-63.
 Journal code: 9Y8; 0372516. ISSN: 0006-291X.
 PUB. COUNTRY: United States
 Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 199906
 ENTRY DATE: Entered STN: 19990714
 Last Updated on STN: 19990714
 Entered Medline: 19990628

AB The PKD1-encoded protein, "**polycystin-1**", has a large N-terminal extracellular portion, multiple transmembrane domains, and a short intracellular C-terminal tail with four tyrosine residues and. . . in kidney development and autosomal dominant polycystic kidney disease (ADPKD) is still unknown. We have subcloned the cDNA encoding the **polycystin-1** C-terminal domain (PKD1-CTD) into a prokaryotic **expression** vector, and site-directed mutagenesis was performed to target the four tyrosine residues and four serine residues in two putative phosphorylation sites. In vitro phosphorylation **assays** were conducted on both wild type and mutant PKD1-CTD fusion proteins. It was found that the wild type PKD1-CTD and. . .

L6 ANSWER 4 OF 4 MEDLINE DUPLICATE 3
 ACCESSION NUMBER: 96202312 MEDLINE
 DOCUMENT NUMBER: 96202312 PubMed ID: 8643665
 TITLE: Polycystin, the polycystic kidney disease 1 protein, is expressed by epithelial cells in fetal, adult, and polycystic kidney.
 AUTHOR: Ward C J; Turley H; Ong A C; Comley M; Biddolph S; Chetty R; Ratcliffe P J; Gattner K; Harris P C
 CORPORATE SOURCE: Medical Research Council Molecular Haematology Unit, Institute of Molecular Medicine, John Radcliffe Hospital, Headington, Oxford, United Kingdom.
 SOURCE: PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (1996 Feb 20) 93 (4) 1524-8.
 Journal code: PV3; 7505876. ISSN: 0027-8424.
 PUB. COUNTRY: United States
 Journal; Article; (JOURNAL ARTICLE)
 LANGUAGE: English
 FILE SEGMENT: Priority Journals
 ENTRY MONTH: 199607
 ENTRY DATE: Entered STN: 19960726
 Last Updated on STN: 19960726
 Entered Medline: 19960717

AB . . . locus of the common genetic disorder autosomal dominant polycystic kidney disease. We have studied PKD1 mRNA, with an RNase protection **assay**, and found widespread **expression** in adult tissue, with high levels in brain and moderate signal in kidney. **Expression** of the PKD1 protein, **polycystin**, was assessed in kidney using monoclonal antibodies to a recombinant protein containing the C terminus of the molecule. In fetal and adult kidney, staining is restricted to epithelial cells. **Expression** in the developing nephron is most prominent in mature tubules, with lesser staining in Bowman's capsule and the proximal ureteric. . . persists in cortical tubules with moderate staining detected in the loops of Henle and collecting ducts. These results suggest that **polycystin**'s major role is in the maintenance of renal epithelial differentiation and organization from early fetal life. Interestingly, **polycystin expression**, monitored at the mRNA level and by immunohistochemistry, appears higher in cystic epithelia, indicating that the disease does not result. . .

=> d his

(FILE 'HOME' ENTERED AT 08:08:46 ON 31 JUL 2001)

FILE 'MEDLINE, BIOSIS, EMBASE, CAPLUS, USPATFULL' ENTERED AT 08:08:58 ON 31 JUL 2001

L1 0 S POLYCYSTIN (P) SCREEN (P) ATPASE (P) COLLAGEN (P) FOCAL (P)
A
L2 12 S POLYCYSTIN (P) SCREEN
L3 4 DUP REM L2 (8 DUPLICATES REMOVED)
L4 1 S POLYCYSTIN (P) ATPASE (P) COLLAGEN (P) FOCAL (P) ADHESION
L5 11 S POLYCYSTIN (P) ASSAY (P) EXPRESSION
L6 4 DUP REM L5 (7 DUPLICATES REMOVED)

=> s polycystin (p) assay (p) atpase (p) focal

L7 1 POLYCYSTIN (P) ASSAY (P) ATPASE (P) FOCAL

=> s polycystin (p) atpase (p) focal

L8 1 POLYCYSTIN (P) ATPASE (P) FOCAL

=> d l8 ibib

L8 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2001 ACS
ACCESSION NUMBER: 2001:507954 CAPLUS
TITLE: Polycystin-based screening methods for compounds
useful in the treatment of polycystic kidney disease
INVENTOR(S): Wilson, Patricia D.; Burrow, Christopher R.
PATENT ASSIGNEE(S): Mount Sinai School of Medicine of New York
University,
USA
SOURCE: PCT Int. Appl., 56 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001050130	A2	20010712	WO 2001-US100317	20010105
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
PRIORITY APPLN. INFO.:			US 2000-478737	A 20000106
			US 2000-689461	A 20001012

=> log y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	52.84	53.05
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION

CA SUBSCRIBER PRICE

-1.76

-1.76

STN INTERNATIONAL LOGOFF AT 08:16:08 ON 31 JUL 2001